

# MSC Attachment Solution

A xeno-free solution that facilitates the attachment and spreading of mesenchymal stem cells cultured in serum-free medium

Cat. No.: 05-752-1H 5 ml 05-752-1F 1 ml Store at: 2-8°C

## Instructions for Use

## **Product Description**

MSC Attachment Solution is a clear sterile solution containing human fibronectin (hFN) obtained by affinity purification on gelatine-sepharose from human plasma. This solution is particularly useful for the culture of cells that are not capable of synthesizing their own biomatrix or when culturing cells in serum-free medium. The solution was qualified for the attachment of human mesenchymal stem cells (hMSC) when culturing in serum-free media.

## Features

- Contains no products of non-human origin.
- Easy-to-use
- Supplied as a ready-to-use stock solution.
- Design for use with serum-free culture system.

## **Precaution and Disclaimer**

- 1. For in vitro diagnostic use.
- 2. Do not use if a visible precipitate is observed in the MSC Attachment Solution.
- 3. Do not use beyond the expiration date indicated on the product label.



### Instruction for use

- Dilute the MSC Attachment Solution 1:100 using sterile DPBS (without Ca++ and Mg++, Cat. No. 02-023-1) and gently mix using a pipette.
- Add the diluted MSC Attachment Solution to the cultureware. Gently agitate the coated cultureawre and verify complete covering of the surface. Use Table 1 for recommended volumes.
- 3. Incubate the coated cultureware for at least 30 minutes in a humidified CO<sub>2</sub> incubator (37°C).

#### 4. Following 30 minutes incubation:

#### For immediate use:

- Gently wash the cultureware with DPBS (For T-25 use at least 5ml).
- Seed cells immediately.
- It is critical that the coating does not dry out.

#### For later use:

- Wrap the coated cultureware with Parafilm<sup>®</sup> and incubate at 2-8°C. Coated cultureware stored under sterile conditions at +2-8°C are stable for 1 week.
- Gently wash the cultureware with DPBS.
- Seed cells immediately.
- It is critical that the coating does not dry out.

#### Table 1. Recommended volume for the coating procedure

Cultureware	Surface area cm <sup>2</sup>	Volume of 1:100 MSC Attachment Solution
96-well	0.34	0.1 ml
24-well	1.9	0.4 ml
12-well	3.9	0.8 ml
6-well / 35 mm ware	9.6	2 ml
6 cm / T 25 Flask	25	5 ml
T75 Flask	75	15 ml

## **Quality Control**

Each lot is tested for sterility as well as for biological performance (the ability to promote the attachment and spreading of MSC in serum-free medium).

The human source material plasma units used in the production of this product have been tested negative for Hepatitis B virus, Hepatitis C virus, (HCV), HIV, HTLV, Parvovirus B-19 by NAT and TPHA.

## **Auxiliary products**

Product name	Cat. No.
MSC NutriStem® XF Basal Medium and Supplement Mix	05-200-1 and 05-201-1
Recombinant Trypsin Solution	03-078-1
Recombinant Trypsin-EDTA Solution	03-079-1
MSC Dissociation Solution	05-075-1
MSC Freezing Solution	05-712-1
Non enzymatic MSC Dissociation Solution	03-077-1
Dulbecco's PBS (w/o Ca & Mg)	02-023-1



BI Biological Industries Culture of Excellence Biological Industries Israel Beit Haemek Ltd. Kibbutz Beit Haemek 25115, Israel T.+972.4.9960595 F.+972.4.9968896 Email: info@bioind.com WWW.bioind.com





CGMP Manufacturing Facility